

The Study of Metamorphic Rocks, Zonation and Isogrades in Garnet Rocks in the Hamadan Area

ZAHRA HOSSEIN MIRZAEI BENI and ZOHREH HOSSEIN MIRZAEI BENI

Young Research Club, Khorasgan (Isfahan) Branch, Islamic Azad University, Isfahan (Iran).

(Received: March 03, 2012; Accepted: April 15, 2012)

ABSTRACT

The study area is a part of the Sanandaj- Sirjan metamorphic belt. We can divide Hamadan metamorphic rocks in three groups: regional metamorphic rocks, contact metamorphic rocks and migmatites. In this area we can't completely divide zonation of contact and regional metamorphic. In some places that contact metamorphic has influenced to low degree regional metamorphic rocks, contact metamorphic zonations are clearly appear, but when contact and regional metamorphic have a same degree or regional metamorphic has high degree than contact metamorphic, we can't distinguish them easily. In Hamadan area regional metamorphic zones are Chlorite± Biotite zone (we haven't garnet rocks in this zone), Biotite± Garnet zone (divided in two sub zone, Biotite and Garnet zone), Andalusite zone, Staurolite zone, Staurolite± Andalusite zone, Sillimanite- Muscovite zone and Sillimanite- Potassium feldspar± Cordierite zone, also contact metamorphic zones are Cordierite zone and Cordierite- Potassium feldspar zone.

Key words: Contact metamorphic; Garnet; Isogrades; Metamorphic zonation; Migmatites; Regional metamorphic.

INTRODUCTION

Garnet crystallizes in cubic system and mostly in dodecahedron (rhomb-dodecahedron) and trapezohedron (tetragon-trioctahedron) crystal forms. General chemical formula of this mineral is: $R_3R'_2(SiO_4)_3$, which bivalent cations (i.e. Mg^{2+} , Fe^{2+} , Mn^{2+} , Ca^{2+}) lie in R site and trivalent cations (i.e. Al^{3+} , Cr^{3+} , Fe^{3+}) in R' site. Commonly, more than one cation lies in R and R' sites and therefore garnet crystals give rise to isomorphous (solid solution) series of minerals. If Al^{3+} is located in R' site, the pyrope group [$(Fe^{2+}, Mg^{2+}, Mn^{2+})_3 Al_2(SiO_4)_3$] with almandine [$(Fe^{2+})_3 Al_2(SiO_4)_3$], pyrope [$(Mg^{2+})_3 Al_2(SiO_4)_3$] and spessartine [$(Mn^{2+})_3 Al_2(SiO_4)_3$] end members will form. If Ca^{2+} is located in R site, the ugrandite group [$(Ca^{2+})_3(Al^{3+}, Fe^{3+}, Cr^{3+})_2(SiO_4)_3$] with grossularite [$Ca_3Al_2(SiO_4)_3$], andradite [$Ca_3(Fe^{3+})_2(SiO_4)_3$] and uvarovite [$Ca_3(Cr^{3+})_2(SiO_4)_3$] end members will form. Some other cations may also be replaced in R and R' sites [1, 2]. The garnet

minerals chemistry in the study area are rich in almandine.

Geological Setting

The study area is a part of the Sanandaj- Sirjan metamorphic belt. The Alvand plutonic complex is the most important plutonic body that regional and contact metamorphic rocks with low to high grade are located around it. The metamorphic sequence comprises pelitic, psammitic, basic, calc-pelitic and calc-silicate rocks. Pelitic rocks are the most abundant lithologies. Pelitic sequence is mostly made up of slates, phillites, micaschists, garnet schists, garnet andalusite (± sillimanite, ± kyanite) schists, garnet staurolite schists, mica hornfelses, garnet hornfelses, garnet andalusite (± fibrolite) hornfelses, cordierite (± andalusite) hornfelses, cordierite K-feldspar hornfelses and sillimanite K-feldspar hornfelses. Major plutonic rocks of this area are granitoids, diorites and gabbroids, which




Fig. 1: Simplified zonation map of the Hamadan area [10]




Figure 2: Mineral assemblage in Garnet zone.




Fig. 3: Mineral assemblage in Chiastolite zone




Fig. 4: Mineral assemblage in Staurolite zone




Fig. 5: Mineral assemblage in Sillimanite muscovite zone




Fig. 6: First mineral assemblage in Sillimanite-potassium feldspar zone




Fig. 7: Second mineral assemblage in Sillimanite- potassium feldspar zone



Fig. 8: Mineral assemblage in Cordierite zone





Fig. 9: Mineral assemblage in Cordierite potassium feldspar zone

Staurolite, Garnet, Biotite, Muscovite, Chlorite, Plagioclase, Graphite and Tourmaline (Figure 4). Their common texture is lepidoporphroblastic with porphyroblasts of garnet, staurolite (up to 15 cm in length).



Sillimanite muscovite zone

Sillimanite andalusite schists contain Quartz, Sillimanite (\pm andalusite), Biotite, Muscovite, Garnet, Plagioclase and Opaque minerals (Figure 5).



Sillimanite- potassium feldspar zone

High grade schists and Migmatites are in this zone. The high grade schists in the regional metamorphic sequence contain Sillimanite, Quartz, Biotite, Muscovite, Garnet, Plagioclase, Potassium feldspar, \pm Andalusite, \pm Kyanite, \pm Staurolite (Figure 6).

Migmatites are a sequence of metatexite-diatexite rocks with various structures such as stromatic, schollen, schlieric and massive. The melanosome mineralogy of the most of the metatexites is very similar to high grade Garnet sillimanite (\pm andalusite and kyanite) schists but Cordierite-bearing interlayers occur, too (Figure 7). Leucosome of migmatites have granoblastic texture and contain Quartz, Plagioclase, Muscovite and \pm Garnet.



Contact metamorphic rocks

Protoliths of the contact metamorphic rocks are similar to those in the regional metamorphic sequence and include abundant metapelitic rocks. Two metamorphic zones are widespread around plutonic bodies.

Cordierite zone


The major rock types in this zone are Cordierite hornfelses. This rocks have porphrogranoblastic texture that containing Quartz, Biotite, Muscovite, contact Cordierite (\pm andalusite), Plagioclase, Garnet, Tourmaline and Opaque

minerals (Figure 8). garnet hornfelses forming at 2.5 ± 0.1 Kbar and $539-569$ °C [3].



Cordierite potassium feldspar zone

The typical mineral assemblage of these rock is Quartz, contact Cordierite (Crd_2), orthoclase, Biotite, minor Plagioclase, Garnet and Opaque minerals (Figure 9).



Minerals assemblage in metamorphic zonation are shown in table 1.

CONCLUSION

We can divide Hamadan metamorphic rocks in three groups: regional metamorphic rocks, contact metamorphic rocks and migmatites. In this area regional metamorphic zones are Chlorite± Biotite zone, Biotite± Garnet zone, Andalusite zone, Staurolite zone, Staurolite± Andalusite zone, Sillimanite- Muscovite zone and Sillimanite-Potassium feldspar± Cordierite zone, also contact metamorphic zones are Cordierite zone and Cordierite- Potassium feldspar zone.

REFERENCES

- Locock, A., An Excel spreadsheet forecast analyses of garnet end-member components, and a synopsis of the crystal chemistry of natural silicate garnets. *Computers and Geosciences*, V, **34**: 1769-1780 (2008).
- Li Li, H., Kuang, X., Mao, A., Li, Y. and Wang, S., Study of local structures and optical spectra for octahedral Fe^{3+} centers in a series of garnet crystals $\text{A}_3\text{B}_2\text{C}_3\text{O}_{12}$ (A = Cd, Ca; B = Al, Ga, Sc, In; C = Ge, Si). *Chemical Physics Letters*, **484**: 387-391 (2010)
- Sepahi, A. A., Whitney D. L. and Baharifar A. A., , Petrogenesis of andalusite-kyanite-sillimanite veins and their host rocks, Sanandaj-Sirjan metamorphic belt, Hamadan, *Iran. J. Met. Geol*, **22**:119-134 (2004).
- Thompson, A.B., A note on the kaolinite-pyrophyllite equilibrium. *Am. J. Sci*, **268**: 454-458 (1970).
- Frey, M., Progressive Low grade metamorphism of a Black Shale Formation, Central Swiss Alps, with special reference to pyrophyllite and margarite bearing assemblages. *J. Petrol*, **19**: 95-135 (1978).
- Whitney, D.L., Mechum, T.A. and Dilek, Y.R., Progressive metamorphism of pelitic rocks from protolith to granulite facies. Dutchess County, New York, USA: Constraints on the timing of fluid infiltration during regional metamorphism. *J. Met. Geol*, **74**: 163-181 (1996).
- Kretz, R., Metamorphic crystallization. John Wiley and Sons, 507 (1994).
- Yang, P. and Pattison, D., Genesis of monazite and Y zoning in garnet from the Black Hills, South Dakota. *J. Lithos*, **88**: 233-253 (2006).
- Norlander, B.H., Whitney, D.L., Teyssier, C. and Vanderhaeghe, O., Partial melting and decompression of the Thor-Odin Dome, Shuswap metamorphic core complex. Canada. *Cord. Lithos*, **61**: 103-125 (2002).
- Sepahi, A.A., Typology and petrogenesis of granitic rocks in the Sanandaj-Sirjan metamorphic belt, Iran: With emphasis on the Alvand plutonic complex. *N. Jb. Geol. Palat. Abn*, **247**: 295-312 (2008).